

UC Agriculture & Natural Resources

Proceedings of the Vertebrate Pest Conference

Title

Bird damage appraisal methods in some agricultural crops

Permalink

<https://escholarship.org/uc/item/32723489>

Journal

Proceedings of the Vertebrate Pest Conference, 6(6)

ISSN

0507-6773

Author

DeHaven, Richard W.

Publication Date

1974

BIRD DAMAGE APPRAISAL METHODS IN SOME AGRICULTURAL CROPS

RICHARD W. DeHAVEN, Wildlife Biologist, U.S. Bureau of Sport Fisheries and Wildlife, Denver Wildlife Research Center Field Station, Davis, California

For more than a decade, personnel of the Section of Bird Damage Control at the Denver Wildlife Research Center have studied agricultural damage by birds. Much of their research has centered on evaluating damage reduction efforts, and in doing this, they have developed and used many damage appraisal methods. This paper outlines briefly those that have been published. It is hoped that these methods will provide other workers with a useful starting point for evaluating bird losses in other test situations or in other surveys over extensive areas. The reader is cautioned, however, that bird damage appraisal is very complex, and in planning sampling designs, the advice of a good agricultural statistician, or at least attention to a good statistical textbook, is important to prevent a wasted effort.

SPROUTING FIELD CROPS

Corn

Damage to fields ranging from 0.8 to 40 acres in size was evaluated by counting normal sprouts and bird-damaged or destroyed sprouts on from 15 to 100 subplots. Subplots usually consisted of a 100-foot section of two adjacent rows (0.016 acre) and were located randomly except in one instance in which they were allocated to field edges and centers at a ratio of 9:1 because bird damage (by pheasants) was stratified. The average number of plants lost per field and the average percentage lost were determined at various intervals after damage began. (Sources: West 1968; West and Dunks 1969; West et al. 1969; Guarino and Forbes 1970; Frank et al. 1970; Stickley and Guarino 1972.)

Rice

Several 1-acre field plots were each gridded into 100 0.01-acre subplots, and a center point for a 1-square-foot circle was randomly chosen within each subplot. Rice seedlings within 6.77 inches of the center point were counted at intervals after damage began and the total plants per field plot estimated. (Source: Besser 1973.)

MATURE FIELD CROPS

Corn

Extensive areas (up to 24 states) were surveyed by examining randomly located fields after the corn had dented (and most damage had stopped). Generally, one to four randomly located subplots, ranging in size from one 15-foot section of a row to one 200-foot section of a row, were selected in each field. (In a recent survey, one 100-foot row was found to be the most efficient.) The number of damaged and undamaged ears were counted, and on damaged ears, the average lengths of damaged and undamaged kernel rows were measured to the nearest 0.1 inch. These lengths were converted to bushels per acre of corn lost by using a mathematically generated table. A bias in the table may somewhat underestimate damage, and this is now being checked. (Sources: De Grazio et al. 1969, Stone et al. 1970, Stone et al. 1971, Stone and Mott 1973.)

Peanuts

Thirty randomly located fields were surveyed, and four subplots, each consisting of a 10-foot-long section of a peanut windrow, were randomly selected in each field. The number of pods opened by birds and the number of pegs where pods had been removed by birds were counted and used to calculate average losses per acre. (Source: Mott et al. 1972.)

Rice

Bird damage to experimental rice was evaluated in relatively small plots (7 x 7, 6 x 6, or 4 x 4 feet). From 10 to 20 rice panicles, clipped 1 inch below the first primary branch, were randomly collected in each plot. Average weight, number of missing or "milked" kernels, and percentage of kernels damaged were calculated. Total plot yield was also determined by harvesting. (Source: DeHaven et al. 1971.)

FRUIT CROPS

Cherries (Control Evaluation)

In randomly selected trees, 50 cherries on the tip of each of eight randomly selected branches were counted and marked off before damage began. Branches were stratified; two were selected for each compass direction (N, S, E, and W), one above and one below the midpoint of the tree. The number of marked cherries damaged or removed by birds and the total percent damage were determined at intervals before harvest. (Source: Guarino *et al.* 1974a,b.)

Cherries (Extensive Survey)

One tree was selected (by a weight randomized scheme) in each of 100 orchards. One terminal branch was randomly selected in each tree, and all green cherries on the branch were counted before damage began. At 3 to 5 days before harvest, all cherries were stripped from the marked branches, the total number of cherries and the number pecked were determined, and these results were compared with predamage counts to calculate percent loss. (Sources: Michigan Crop Reporting Service 1972; Stone 1974.)

Blueberries

Blueberry bushes (selected randomly in one test and along a line transect in another) were chosen for subsampling in plantings ranging from 0.5 to 8 acres in size. Damage estimates were based on percentage losses (by actual count) to groups of 10 ripe berries on each bush. In one test, berry groups were selected and marked before damage began; in the second, damage had started, and marked branches were thinned to 10 undamaged berries. (Sources: Bollengier *et al.* 1974, Stone *et al.* 1974.)

LITERATURE CITED

- BESSER, J. F. 1973. Protecting seeded rice from blackbirds with methiocarb. *Int. Rice Comm. Newsl.* 22(3):9-14.
- BOLLENGIER, R. M., J. L. GUARINO, and C. P. STONE. 1974. Aerially applied methiocarb spray for protecting wild lowbush blueberries from birds. *Proc. 6th Bird Control Seminar*, Bowling Green State University, Bowling Green, Ohio. (in press)
- DE GRAZIO, J. W., J. F. BESSER, J. L. GUARINO, C. M. LOVELESS, and J. L. OLDEMEYER. 1969. A method for appraising blackbird damage to corn. *J. Wildl. Manage.* 33(4):988-994.
- DeHAVEN, R. W., J. L. GUARINO, F. T. CRASE, and E. W. SCHAFER, JR. 1971. Methiocarb for repelling blackbirds from ripening rice. *Int. Rice Comm. Newsl.* 20(4):25-30.
- FRANK, F. E., E. W. SCHAFER, JR., and J. L. GUARINO. 1970. Laboratory and field studies with an avian repellent for sprouting seeds. *Proc. 5th Bird Control Seminar*, Bowling Green State Univ., Bowling Green, Ohio. pp. 86-89.
- GUARINO, J. L., and J. E. FORBES. 1970. Preventing bird damage to sprouting corn with a carbamate repellent. *New York Fish and Game J.* 17(2):117-120.
- _____, W. F. SHAKE, and E. W. SCHAFER, JR. 1974a. Reducing bird damage to ripening cherries with methiocarb. *J. Wildl. Manage.* (in press)
- _____, C. P. STONE, and W. F. SHAKE. 1974b. A low-level treatment of the avian repellent, methiocarb, on ripening sweet cherries. *Proc. 6th Bird Control Seminar*, Bowling Green State Univ., Bowling Green, Ohio. (in press)
- MICHIGAN CROP REPORTING SERVICE. 1972. Enumerators manual: 1972 tart cherry objective yield survey. U. S. Dep. Agric. and Michigan Dep. Agric., Lansing. 73pp.
- MOTT, D. F., J. F. BESSER, R. R. WEST, and J. W. DE GRAZIO. 1972. Bird damage to peanuts and methods for alleviating the problem. *Proc. 5th Vertebrate Pest Conf.*, Fresno, Calif. pp. 118-120.
- STICKLEY, A. R., JR., and J. L. GUARINO. 1972. A repellent for protecting corn seed from blackbirds and crows. *J. Wildl. Manage.* 36(1):150-152.
- STONE, C. P. 1974. Bird damage to tart cherries in Michigan, 1972. *Proc. 6th Bird Control Seminar*, Bowling Green State Univ., Bowling Green, Ohio. (in press)
- _____, and D. F. MOTT. 1973. Bird damage to ripening field corn in the United States, 1971. U. S. Dep. Int. Bur. Sport Fisheries and Wildl. *Wildl. Leaf.* 505. 8pp.
- _____, D. F. MOTT, J. F. BESSER, and J. W. DE GRAZIO. 1972. Bird damage to corn in the United States in 1970. *Wilson Bull.* 84(1):101-105.
- _____, D. F. MOTT, J. E. FORBES, and C. P. BREIDENSTEIN. 1973. Bird damage to field corn in New York in 1971. *New York Fish and Game J.* 20(1):68-73.
- _____, W. F. SHAKE, and D. J. LANGOWSKI. 1974. Reducing bird damage to highbush blueberries in Michigan with a carbamate repellent. *Wildl. Soc. Bull.* (in press)

WEST, R. R. 1968. Protecting sprouting corn from pheasants with DRC-736. Trans. 13th Ann. Conf. Central Mountain & Plains Sect., Wildl. Soc. p 12.

_____, R. B. BRUNTON, and D. J. CUNNINGHAM. 1969. Repelling pheasants from sprouting corn with a carbamate insecticide. J. Wildl. Manage. 33(1):216-219.

_____, and J. H. DUNKS. 1969. Repelling boat-tailed grackles from sprouting corn with a carbamate compound. Texas J. Sci. 21(2):231-233.